arXiv:1209.6013v1 [cs.FL] 26 Sep 2012

Abelian Repetitions in Sturmian Words

Gabriele Fici[1], Alessio Langiu[2], Thierry Lecroq[3], Arnaud Lefebvre[3], Filippo Mignosi[4],
and Élise Prieur-Gaston[3]

[1] I3S, CNRS & Université Nice Sophia Antipolis, France
Gabriele.Fici@unice.fr
[2] Department of Informatics, King's College London, UK
Alessio.Langiu@kcl.ac.uk
[3] Université de Rouen, LITIS EA4108, 76821 Mont-Saint-Aignan Cedex, France
{Thierry.Lecroq,Arnaud.Lefebvre,Elise.Prieur}@univ-rouen.fr
[4] Dipartimento di Informatica, Università dell'Aquila, L'Aquila, Italy
Filippo.Mignosi@di.univaq.it

Abstract. We investigate abelian repetitions in Sturmian words. We exploit a bijection between factors of Sturmian words and subintervals of the unitary segment that allows us to study the periods of abelian repetitions by using classical results of elementary Number Theory. We prove that in any Sturmian word the ratio between the maximal exponent of an abelian repetition of period m and m converges to a number greater than or equal to $\sqrt{5}$, and the equality holds for the Fibonacci infinite word. We further prove that the longest prefix of the Fibonacci infinite word that is an abelian repetition of period F_j, $j > 1$, has length $F_j(F_{j+1} + F_{j-1} + 1) - 2$ if j is even or $F_j(F_{j+1} + F_{j-1}) - 2$ if j is odd. This allows us to give an exact formula for the smallest abelian periods of the Fibonacci finite words. More precisely, we prove that for $j \geq 3$, the Fibonacci word f_j has abelian period equal to F_n, where $n = \lfloor j/2 \rfloor$ if $j = 0, 1, 2 \mod 4$, or $n = 1 + \lfloor j/2 \rfloor$ if $j = 3 \mod 4$.

Keywords: Formal Languages, Combinatorics on Words, Stringology, abelian repetitions, Sturmian words, Fibonacci words.

1 Introduction

The study of repetitions in words is a classical subject in Theoretical Computer Science both from the combinatorial and the algorithmic point of view. Repetitions are strictly related to the notion of periodicity. Recall that a word w of length $|w|$ has a *period* $p > 0$ if $w[i] = w[i+p]$ for any $1 \leqslant i \leqslant |w|-p$, where $w[i]$ is the symbol in position i of w. Every word w has a minimal period $p \leq |w|$. If $|w|/p \geq 1$, w is called a *repetition* of period p and exponent $|w|/p$. When $|w|/p = k$ is an integer, the word w is called an *integer power*, since it can be written as $w = u^k$, that is, w is the concatenation of k copies of a word u of length p. If instead $|w|/p$ is not an integer, the word w is called a *fractional power*. So one can write $w = u^k v$, where v is the prefix of u such that $|w|/p = k + |v|/|u|$. For example, the word $w = aabaaba$ is a 7/3-power since it has minimal period 3 and length 7. A classical reference on periodicity is [29, Chap. 7].

Abelian properties concerning words have been studied since the very beginning of Formal Languages and Combinatorics on Words. The notion of Parikh vector has become a *standard* and it is often used without an explicit reference to the 1966 Parikh paper [35], that is however cited more than 450 times. On the other hand abelian powers were first considered in 1961 by Erdös [20] as a natural generalization of usual powers. Researches concerning abelian properties of words and languages never stopped and they developed in different directions (see for instance [9, 12, 17, 18, 23, 26, 30, 37, 39, 40, 41]). There is a recent increasing of interest on abelian properties linked to periodicity (see [2, 3, 6, 7, 10, 11, 16, 19, 23, 39, 40, 42] and references therein).

Recall that the Parikh vector $\mathcal{P}_w$ of a finite word w enumerates the cardinality of each letter of the alphabet in w. For example, given the alphabet $\Sigma = \{a, b, c\}$, the Parikh vector of the word $w = aaba$ is $\mathcal{P}_w = (3, 1, 0)$. Roughly speaking, two words have the same Parikh vector if one can be obtained from the other by permuting letters.

We say that the word w is an *abelian repetition* of (abelian) period m and exponent $|w|/m$ if w can be written as $w = u_0u_1 \cdots u_{k-1}u_k$ for words u_i and an integer $k > 2$, where for $0 < i < k$ all the u_i's have the same Parikh vector whose sum of components is m and the Parikh vectors of u_0 and u_k are contained in $\mathcal{P}$ (see [13]). When u_0 and u_k are empty, w is called an *abelian power* or *weak repetition* [15]. For example, the word $w = abaab$ is an abelian repetition of period 2, since one can set $u_0 = a$, $u_1 = ba$, $u_2 = ab$ and $u_3 = \varepsilon$, where ε denotes the empty word.

It is well known that Sturmian words and Fibonacci words, in particular, are extremal cases for several problems related to repetitions (see for example [14, 24, 34]) and are worst-case examples for classical pattern matching algorithms, e.g. Knuth-Morris-Pratt [1, 28]. There exists a huge bibliography concerning Sturmian words (see for instance the survey papers [4, 5, 27], [29, Chap. 3], [38, Chap. 6] and references therein). In particular, there is an analogous result to the one presented in this paper concerning classical repetitions in the Fibonacci infinite word [33]. This result was subsequently extended to the class of Sturmian words in several other papers. In [31], a bijection between factors of Sturmian words and subintervals of the unitary segment is described. In this paper we show that this bijection preserves abelian properties of factors (see Proposition 4). Therefore we are able to apply techniques of Number Theory coupled with Combinatorics on Words to obtain our main results. More precisely, we prove that in any Sturmian word the ratio between the maximal exponent of an abelian repetition of period m and m converges to a number greater than or equal to $\sqrt{5}$, and the equality holds for the Fibonacci infinite word. We further prove that the longest prefix of the Fibonacci infinite word that is an abelian repetition of period F_j, $j > 1$, has length $F_j(F_{j+1} + F_{j-1} + 1) - 2$ if j is even or $F_j(F_{j+1} + F_{j-1}) - 2$ if j is odd. This allows us to give an exact formula for the smallest abelian periods of the Fibonacci finite words. More precisely, we prove that for $j \geq 3$, the Fibonacci word f_j has abelian period equal to F_n, where $n = \lfloor j/2 \rfloor$ if $j = 0, 1, 2 \mod 4$, or $n = 1 + \lfloor j/2 \rfloor$ if $j = 3 \mod 4$.

2 Preliminaries

Let $\Sigma = \{a_1, a_2, \ldots, a_\sigma\}$ be a finite ordered alphabet of cardinality σ and Σ^* the set of words over Σ. We denote by $|w|$ the length of the word w. We write $w[i]$ the i-th symbol of w and $w[i\,.\,.\,j]$ the factor of w from the i-th symbol to the j-th symbol, with $1 \leqslant i \leqslant j \leqslant |w|$. We denote by $|w|_a$ the number of occurrences of the symbol $a \in \Sigma$ in the word w.

The *Parikh vector* of a word w, denoted by $\mathcal{P}_w$, counts the occurrences of each letter of Σ in w, that is, $\mathcal{P}_w = (|w|_{a_1}, \ldots, |w|_{a_\sigma})$. Given the Parikh vector $\mathcal{P}_w$ of a word w, we denote by $\mathcal{P}_w[i]$ its i-th component and by $|\mathcal{P}_w|$ the sum of its components. Thus, for a word w and $1 \leqslant i \leqslant \sigma$, we have $\mathcal{P}_w[i] = |w|_{a_i}$ and $|\mathcal{P}_w| = \sum_{i=1}^{\sigma} \mathcal{P}_w[i] = |w|$. Finally, given two Parikh vectors $\mathcal{P}, \mathcal{Q}$, we write $\mathcal{P} \subset \mathcal{Q}$ if $\mathcal{P}[i] \leqslant \mathcal{Q}[i]$ for every $1 \leqslant i \leqslant \sigma$ and $|\mathcal{P}| < |\mathcal{Q}|$.

Following [13], we give the definition below.

Definition 1. *A word w is an abelian repetition of period $m > 0$ and exponent $|w|/m$ if one can write $w = u_0u_1 \cdots u_{k-1}u_k$ for some $k > 2$ such that $\mathcal{P}_{u_0} \subset \mathcal{P}_{u_1} = \cdots = \mathcal{P}_{u_{k-1}} \supset \mathcal{P}_{u_k}$, and $|\mathcal{P}_{u_1}| = \ldots = |\mathcal{P}_{u_{k-1}}| = m$. An abelian power is an abelian repetition in which $u_0 = u_k = \varepsilon$.*

We call u_0 and u_k respectively the *head* and the *tail* of the abelian repetition. Notice that the length $t = |u_k|$ of the tail is uniquely determined by h, m and $|w|$, namely $t = (|w| - h) \bmod m$.

In the rest of the paper, when we refer to an abelian repetition of period m, we always suppose that m is the minimal abelian period of w.

Example 1. The word $w = abaababa$ is an abelian repetition of period 2, since one can write $w = a \cdot ba \cdot ab \cdot ab \cdot a$. Notice that w is also an abelian repetition of period 3, since $w = aba \cdot aba \cdot ba$.

2.1 Sturmian words

Here and in the rest of the paper, we fix the alphabet $\Sigma = \{\mathsf{a,b}\}$. We start by recalling a bijection between factors of Sturmian words and subintervals of the unitary segment introduced in [31].

Let α and ρ be two real numbers with $\alpha \in (0, 1)$. Following the notations of [22], the fractional part of a number r is defined by $\{r\} = r - \lfloor r \rfloor$, where $\lfloor r \rfloor$ is the greatest integer smaller than or equal to r. Therefore, for $\alpha \in (0, 1)$, one has that $\{-\alpha\} = 1 - \alpha$.

The sequence $\{n\alpha + \rho\}, n > 0$, defines an infinite word $s_{\alpha,\rho} = a_1(\alpha, \rho)a_2(\alpha, \rho) \cdots$ by the rule

$$a_n(\alpha, \rho) = \begin{cases} \mathsf{b} \text{ if } \{n\alpha + \rho\} \in [0, \{-\alpha\}), \\ \mathsf{a} \text{ if } \{n\alpha + \rho\} \in [\{-\alpha\}, 1). \end{cases}$$

See Fig. 1 for a graphical illustration.

We shall write a_n instead of $a_n(\alpha, \rho)$ whenever there is no possibility of mistake. If $\alpha = p/q$, with p and q coprime integers, then it is easy to prove that the word $s_{\alpha,\rho}$ is periodic and q is its minimal period. If instead α is irrational, then $s_{\alpha,\rho}$ is not periodic and is called a *Sturmian word*. Therefore, in the rest of the paper, we always suppose α irrational.

Example 2. For $\alpha = \phi - 1$ and $\rho = 0$, where $\phi = (1 + \sqrt{5})/2$ is the golden ratio, one obtains the Fibonacci infinite word

$$f = s_{\alpha,0} = \mathsf{abaababaabaababaababa} \cdots$$

Remark 1. Since $\alpha \in (0, 1)$, we have $\{-i\alpha\} \neq \{-(i+1)\alpha\}$ for any natural number i. We shall use this fact freely and with no explicit mention.

It is possible to prove (see [31, Corollary 2.3]) that the following proposition holds.

Proposition 1. *Let α and ρ be real numbers, with $\alpha \in (0, 1)$ irrational. For any natural numbers n, i, with $n > 0$, if $\{-(i + 1)\alpha\} < \{-i\alpha\}$ then*

$$a_{n+i} = \mathsf{a} \iff \{n\alpha + \rho\} \in [\{-(i + 1)\alpha\}, \{-i\alpha\}),$$

whereas if $\{-i\alpha\} < \{-(i + 1)\alpha\})$ then

$$a_{n+i} = \mathsf{a} \iff \{n\alpha + \rho\} \in [0, \{-i\alpha\}) \cup [\{-(i + 1)\alpha\}, 1).$$



Fig. 1. An application of Proposition 1 when $\alpha = \phi - 1 \approx 0.618$ (thus $\{-\alpha\} \approx 0.382$) for $i = 0$. If $\{n\alpha + \rho\} \in [\{-\alpha\}, 1)$, then $a_n = \mathsf{a}$; otherwise $a_n = \mathsf{b}$.



Fig. 2. An application of Proposition 1 when $\alpha = \phi - 1 \approx 0.618$ (thus $\{-\alpha\} \approx 0.382$) for $i = 1$. If $\{n\alpha + \rho\} \in [0, \{-\alpha\}) \cup [\{-2\alpha\}, 1)$, then $a_{n+1} = \mathsf{a}$; otherwise $a_{n+1} = \mathsf{b}$.



Fig. 3. A single graphic representation of the information given in Fig. 1 and 2. If $\{n\alpha + \rho\} \in [0, \{-\alpha\}) = L_0(\alpha, 2)$, then $a_n = \mathsf{b}$, $a_{n+1} = \mathsf{a}$. If $\{n\alpha + \rho\} \in [\{-\alpha\}, \{-2\alpha\}) = L_1(\alpha, 2)$, then $a_n = \mathsf{a}$, $a_{n+1} = \mathsf{b}$. If $\{n\alpha + \rho\} \in [\{-2\alpha\}, 1) = L_2(\alpha, 2)$, then $a_n = \mathsf{a}$, $a_{n+1} = \mathsf{a}$.

In Fig. 1 and 2 we display a graphical representation of the formula given in Proposition 1 for $\alpha = \phi - 1$ when $i = 0$ and $i = 1$, respectively. In Fig. 3 we present within a single graphic the situations illustrated in Fig. 1 and 2.

Let m be a positive integer. Consider the $m + 1$ points $0, 1, \{-i\alpha\}$, for $0 < i \leq m$. Rearranging these points in increasing order one has:

$$0 = c_0(\alpha, m) < c_1(\alpha, m) < \cdots < c_k(\alpha, m) < \cdots < c_m(\alpha, m) < c_{m+1}(\alpha, m) = 1.$$

One can also define the non-empty intervals $L_k(\alpha, m) = [c_k(\alpha, m), c_{k+1}(\alpha, m)),\ 0 \leq k \leq m$.

By using Proposition 1, it is possible to associate with each interval $L_k(\alpha, m)$ a factor of length m of the word $s_{\alpha,\rho}$, and this correspondence is bijective (see [32]).

Proposition 2. *Each factor of $s_{\alpha,\rho}$ of length m, $a_n a_{n+1} \ldots a_{n+m-1}$, depends only on the interval $L_k(\alpha, m)$ containing the point $\{n\alpha + \rho\}$; more precisely, it depends only on the set $I_k(\alpha, m)$ of integers $i \in \{0, 1, \ldots, m-1\}$ such that either $\{-(i+1)\alpha\} < \{-i\alpha\}$ and $c_k(\alpha, m) \in [\{-(i+1)\alpha\}, \{-i\alpha\})$ or $\{-(i+1)\alpha\} > \{-i\alpha\}$ and $c_k(\alpha, m) \notin [\{-i\alpha\}, \{-(i+1)\alpha\})$. The set $I_k(\alpha, m)$ is the set of the integers i, $0 \leq i \leq m-1$, such that $a_{n+i} = \mathsf{a}$.*

Corollary 1. *Since the set of factors of $s_{\alpha,\rho}$ depends only on the sequence $\{-i\alpha\}$, $i > 0$, it does not depend on ρ.*

Definition 2. *We call the bijective correspondence in Proposition 2 the* Sturmian bijection.

Example 3. Let $\alpha = \phi - 1$. In Fig. 3 we show an example of the Sturmian bijection when $m = 2$. The ordered sequence of points defining the intervals $L_k(\alpha, 2)$ is

$$c_0(\alpha,2) = 0,\ c_1(\alpha,2) = \{-\alpha\} \approx 0.382,\ c_2(\alpha,2) = \{-2\alpha\} \approx 0.764,\ c_3(\alpha,2) = 1.$$

In Fig. 4 we show an example of the Sturmian bijection when $m = 6$. Below each interval there is the factor of s_α of length $m = 6$ associated with that interval. The prefix of length 6 of the Fibonacci word corresponds to the factor below the interval containing α (so, for $n = 1$ and $\rho = 0$). Notice that all the factors of length 6 of the Fibonacci word appear, and moreover they appear in lexicographic order from right to left. This property concerning lexicographic order holds for any Sturmian word and any length m of factors and it is proved in next proposition.

Next proposition is not strictly linked to main results of this paper, but it is of independent interest and it is related to recent researches on Sturmian words (see [8,21, 25,36]).

Proposition 3. *For any $m \geq 1$ we have that $k < k'$ if and only if the factor $t_{\alpha,\rho,m}$ associated to $L_k(\alpha, m)$ in the Sturmian bijection is greater than the factor $t'_{\alpha,\rho,m}$ associated to $L_{k'}(\alpha, m)$, following the lexicographic order where $\mathsf{a} < \mathsf{b}$.*

Proof. The proof is by induction on m. The base of the induction is true by very definition of the sequence of the two intervals $L_i(\alpha, 1)$, $i = 0, 1$ and the case $\alpha = \phi - 1$, is showed in Figure 1. Suppose that the property holds for m and let us prove it for $m+1$. The sequence of intervals $L_i(\alpha, m)$, $i = 0, \ldots, m$ is the same sequence of intervals $L_i(\alpha, m+1)$, $i = 0, \ldots, m+1$ except for the interval $L_{\hat{k}}(\alpha, m)$ containing $\{-(m+1)\alpha\}$ that is subdivided into two intervals in the second sequence. Therefore the prefixes of length m of the two factors $t_{\alpha,\rho,m+1}$ and $t'_{\alpha,\rho,m+1}$ are equal if and only if $k' = k+1$ and $L_{\hat{k}}(\alpha, m) = L_k(\alpha, m+1) \cup L'_k(\alpha, m+1)$. In this case the last letter of $t_{\alpha,\rho,m+1}$ is b and the last letter of $t'_{\alpha,\rho,m+1}$ is a and the statement of this proposition is verified. Otherwise we use the induction on the prefix of length m of both factors. ☐

In this paper, we present a new semantic property of the Sturmian bijection, which will be described in the next section, that will allow us to use some Number Theory techniques to deal with abelian repetitions in Sturmian words and, in particular, in the Fibonacci word. Similar techniques were used in [40] to derive nice results on abelian powers on Sturmian words, that are repetitions with empty head and tail.

3 On the Sturmian bijection and Parikh vectors

Let $s_{\alpha,\rho}$ be a Sturmian word. Since we are mainly interested in the set of factors of $s_{\alpha,\rho}$, we do not lose generality, by Corollary 1, supposing $\rho = 0$. For simplicity of notation, we will therefore write s_α instead of $s_{\alpha,0}$.

In this section, we describe some properties of the Sturmian bijection between the factors of length m of s_α and the intervals $L_k(\alpha, m)$ that we will use in the rest of the paper.

Proposition 4. *Under the Sturmian bijection, all the factors corresponding to an interval $c_k(\alpha, m) = [x, y)$ with $x \geq \{-m\alpha\}$ have the same Parikh vector $v_1(\alpha, m)$ and all the factors corresponding to an interval $[x, y)$ with $y \leq \{-m\alpha\}$ have the same Parikh vector $v_2(\alpha, m)$. Moreover, one has $v_1(\alpha, m)[1] = v_2(\alpha, m)[1] + 1$.*


{−3α} {−6α}{−α} {−4α} {−2α} {−5α}
c0(α, 6) c1(α, 6) c2(α, 6)c3(α, 6) c4(α, 6) α c5(α, 6) c6(α, 6)c7(α, 6)
0 0.145... 0.291...0.381... 0.527... 0.763... 0.909... 1
(a_n) (a_{n+1}) (a_{n+2}) (a_{n+3}) (a_{n+4}) (a_{n+5})
b a b a a b
b a a b a b
b a a b a a
a b a b a a
a b a a b a
a a b a b a
a a b a a b

Fig. 4. Subintervals $L_k(\alpha, m)$ obtained for $\alpha = \phi - 1$ and $m = 6$. Below each interval there is the factor of s_α of length 6 associated to that interval. For $\rho = 0$ and $n = 1$, the prefix of length 6 of the Fibonacci word is associated to $L_4(\alpha, 6) = [c_4(\alpha, 6), c_5(\alpha, 6))$, that is the interval containing α.

Proof. We prove the statement by induction on m. The case $m = 1$ is true by definition. Suppose the statement true for m and let us prove it for $m + 1$. We have two cases:

Case 1) $\{-(m + 1)\alpha\} < \{-m\alpha\}$;

Case 2) $\{-m\alpha\} < \{-(m + 1)\alpha\}$.

We give the proof for the first case since the proof for the second case is analogous. By Proposition 2, using the same notation, if $c_k(\alpha, m + 1) \in [\{-(m + 1)\alpha\}, \{-m\alpha\})$ then $a_{n+m} = \mathsf{a}$ otherwise $a_{n+m} = \mathsf{b}$. If $c_k(\alpha, m + 1) \in [\{-(m + 1)\alpha\}, \{-m\alpha\})$ we also have that $L_k(\alpha, m + 1) = [c_k(\alpha, m + 1), c_{k+1}(\alpha, m + 1)) = [c_{k-1}(\alpha, m), c_k(\alpha, m))$ except in the case $c_k(\alpha, m+1) = \{-(m+1)\alpha\}$ where $L_k(\alpha, m+1) = [\{-(m+1)\alpha\}, c_k(\alpha, m)) \subset [c_{k-1}(\alpha, m), c_k(\alpha, m))$. This is because in order to obtain the sequence $c_k(\alpha, m + 1)$ we add the point $\{-(m + 1)\alpha\}$ to the sequence $c_k(\alpha, m)$, and, consequently, the index of all points in the original sequence greater than $\{-(m + 1)\alpha\}$ is increased by one in the newly obtained sequence. Therefore, the Parikh vector of the factors corresponding to the intervals $L_k(\alpha, m + 1)$, where $c_k(\alpha, m + 1) \in [\{-(m + 1)\alpha\}, \{-m\alpha\})$, is equal to $v_2(\alpha, m) + (1, 0)$, since $a_{m+1} = \mathsf{a}$. For the other intervals, since $a_{m+1} = \mathsf{b}$, the Parikh vector of the factors corresponding to the intervals is equal to $v_2(\alpha, m) + (0, 1)$ if $c_k(\alpha, m + 1) < \{-(m + 1)\alpha\} < \{-m\alpha\}$ or to $v_1(\alpha, m) + (0, 1)$ if $c_k(\alpha, m + 1) \geq \{-m\alpha\}$.

Putting all together, we have that $v_2(\alpha, m+1) = v_2(\alpha, m)+(0, 1)$ while $v_1(\alpha, m+1) = v_2(\alpha, m)+(1, 0) = v_1(\alpha, m)+(0, 1)$, and, consequently, $v_1(\alpha, m)[1] = v_2(\alpha, m)[1]+1$. □

The reader can see in Fig. 4 that the factors of length 6 corresponding to an interval to the left of $\{-6(\phi - 1)\}$ have Parikh vector $(3, 3)$, while the others have Parikh vector $(4, 2)$.

We now address the following questions:

1. Given m, how long can be the longest abelian power of period m in s_α?
2. Given m, how long can be the longest abelian power of period m that is a prefix of s_α?
3. What can we say in the particular case of the Fibonacci word?

We start with a corollary, whose proof comes straightforwardly from Proposition 4.

Corollary 2. *Let w be an abelian power of period m and exponent $k + 1$ appearing in s_α in position n. Then all the points in the sequence $\{n\alpha\}, \{(n + m)\alpha\}, \{(n +$*

$2m)\alpha\}, \ldots, \{(n+km)\alpha\}$ are in the same subinterval in which $[0,1)$ is subdivided by the point $\{-m\alpha\}$, that is, either $[0,\{-m\alpha\})$ or $[\{-m\alpha\},1)$.

Proposition 5. *If $k \geq 1$, the $k+1$ points of Corollary 2 are naturally ordered. That is, if $\{m\alpha\} < 0.5$, then they are all in the subinterval $[0,\{-m\alpha\})$ and one has $\{n\alpha\} < \{(n+m)\alpha\} < \cdots < \{(n+km)\alpha\}$; if $\{m\alpha\} > 0.5$ then they are all in the interval $[\{-m\alpha\},1)$ and one has $\{(n+km)\alpha\} < \{(n+(k-1)m)\alpha\} < \cdots < \{n\alpha\}$.*

Theorem 1. *Let m be a positive integer such that $\{m\alpha\} < 0.5$ (resp. $\{m\alpha\} > 0.5$). Then:*

1. *In s_α there is an abelian power of period m and exponent $k \geq 2$ if and only if $\{m\alpha\} < \frac{1}{k}$ (resp. $\{-m\alpha\} < \frac{1}{k}$).*
2. *If in s_α there is an abelian power of period m and exponent $k \geq 2$ starting in position i with $\{i\alpha\} \geq \{m\alpha\}$ (resp. $\{i\alpha\} \leq \{m\alpha\}$), then $\{m\alpha\} < \frac{1}{k+1}$ (resp. $\{-m\alpha\} < \frac{1}{k+1}$). Conversely, if $\{m\alpha\} < \frac{1}{k+1}$ (resp. $\{-m\alpha\} < \frac{1}{k+1}$), then there is an abelian power of period m and exponent $k \geq 2$ starting in position m.*

Proof. (Sketch) Let m be a positive integer such that $\{m\alpha\} < 0.5$. If in s_α there is an abelian power of period m and exponent $k \geq 2$, then by previous proposition, all the k points of Corollary 2 are naturally ordered and they must lie in the interval $[0,\{-m\alpha\})$. The distance between any two consecutive such points is $\{m\alpha\}$. Therefore, the size of this interval, that is equal to $1-\{m\alpha\}$, must be smaller than $(k-1)\{m\alpha\}$. From this relation, one derives that $\{m\alpha\} < 1/k$.

Conversely, suppose that $\{m\alpha\} < 1/k$. Since α is irrational, the sequence $\{i\alpha\}$ is dense in $[0,1)$ (see [22, Chap. XXIII]). Therefore, one can find a number i close to 0 such that the points of Corollary 2 all lie in the interval $[0,\{-m\alpha\})$. The thesis follows by Proposition 4. The case $\{m\alpha\} > 0.5$ can be proved in an analogous way.

Also the second part of the theorem is dealt in a similar way. The only difference is that the first point of the sequence cannot be close to 0 but has to be far away at least $\{-m\alpha\}$, and this fact leads to the stronger condition $\{m\alpha\} < 1/(k+1)$. □

The previous theorem allows us to deal with abelian repetitions in a Sturmian word s_α by using classical results on the approximation of the irrational α by rationals. This is a classical topic in Number Theory. Since the number $\phi - 1$ has special properties within this topic, we have in turn specific results for the Fibonacci word.

4 Approximating irrationals by rationals and abelian repetitions

We recall some classical results of Number Theory. For any notation non explicitly defined in this section we refer to [22, Chap. X, XI].

The sequence $F_0 = 1, F_1 = 1, F_{j+1} = F_j + F_{j-1}$ for $j \geq 1$ is the well known sequence of Fibonacci numbers. The sequence of fractions $\frac{F_{j+1}}{F_j}$ converge to $\phi = \frac{\sqrt{5}+1}{2}$, while the sequence $\frac{F_j}{F_{j+1}}$ converges to $\phi - 1 = \frac{\sqrt{5}-1}{2}$. Moreover, the sequences $\frac{F_{j+1}}{F_j}$ and $0 = \frac{0}{1}$, $\frac{F_j}{F_{j+1}}$, $j = 0, 1, \ldots$, are the sequences of convergents, in the development in continued fractions, of ϕ and $\phi - 1$ respectively.

Concerning the approximation given by the above convergents, the following result holds (see [22, Chap. X, Theorem 171] and [22, Chap. XI, Section 11.8]).

Theorem 2. *For any $j > 0$,*

$$\phi - \frac{F_{j+1}}{F_j} = (\phi - 1) - \frac{F_{j-1}}{F_j} = \frac{(-1)^j}{F_j(\phi F_j + F_{j-1})}.$$

We also report the following theorems (see [22, Chap. XI, Theorem 193] and the proof of [22, Chap. XI, Theorem 194]).

Theorem 3. *Any irrational α has an infinity of approximations which satisfy*

$$\left|\frac{p}{q} - \alpha\right| < \frac{1}{\sqrt{5}q^2}.$$

Theorem 4. *Suppose $\alpha = \phi - 1$. If $A > \sqrt{5}$, the inequality*

$$\left|\frac{p}{q} - \alpha\right| < \frac{1}{Aq^2}$$

has only a finite number of solutions.

The last two theorems coupled with the first part of Theorem 1 allow us to derive the next result.

Theorem 5. *Let s_α be a Sturmian word. For any integer $q > 1$, let k_q be the maximal exponent of an abelian repetition of period q in s_α. Then*

$$\limsup \frac{k_q}{q} \geq \sqrt{5}$$

and the equality holds when $\alpha = \phi - 1$.

Proof. (Sketch) Pick an approximation p/q of α such that $|p/q - \alpha| < 1/\sqrt{5}q^2$. Suppose that $p/q - \alpha < 0$, the proof when $p/q - \alpha > 0$ is analogous.

We have that $q\alpha - p < 1/\sqrt{5}q \leq 0.5$. Hence, $\{q\alpha\} < 1/\sqrt{5}q$ and, by Theorem 1, there is in s_α an abelian repetition of exponent $\sqrt{5}q$ and period q.

If $\limsup k_q/q = A > \sqrt{5}$ then, by Theorem 1, the inequality $|p/q - \alpha| < 1/Aq^2$ has infinitely many solutions. This is impossible by Theorem 4. ☐

We now study the abelian repetitions that are prefixes of the Fibonacci infinite word. For this, we will make use of the second part of Theorem 1.

Notice that an abelian repetition of period m appearing as a prefix of the Fibonacci word can have a head of length equal to $m - 1$ at most. Therefore, we have to check all the abelian powers that start in position i for every $i = 1, \ldots, m$.

In order to do this, we report here another result (see [22, Chap. X, Theorem 182]).

Theorem 6. *Let p_n/q_n be the n-th convergent to α. If $n > 1$, $0 < q \leq q_n$ and $p/q \neq p_n/q_n$, then $|p_n - q_n\alpha| < |p - q\alpha|$.*

We translate the statement of the theorem above in terms more convenient to our setting.

Corollary 3. *Suppose that $m > 1$ is the denominator of a convergent to α and that $\{m\alpha\} < 0.5$ (resp. $\{m\alpha\} > 0.5$). For any i, $1 \leq i < m$, one has $\{i\alpha\} \geq \{m\alpha\}$ (resp. $\{i\alpha\} \leq \{m\alpha\}$).*

From the previous corollary, we have that if $m > 1$ is a Fibonacci number, for $\alpha = \phi - 1$ the hypotheses of the second part of Theorem 1 are satisfied. The next proposition is a direct consequence of the previous corollary, Theorem 1 and Theorem 2.

Proposition 6. *Let $j > 1$. In the Fibonacci infinite word, the longest abelian power having period F_j and starting in a position $i \leq F_j$ has an occurrence starting in position F_j and has exponent*

$$\lfloor \phi F_j + F_{j-1} \rfloor - 1 = \begin{cases} F_{j+1} + F_{j-1} - 1 & \textit{if } j \textit{ is even;} \\ F_{j+1} + F_{j-1} - 2 & \textit{if } j \textit{ is odd.} \end{cases}$$

The following theorem provides a formula for computing the length of the longest abelian repetition appearing as a prefix in the Fibonacci infinite word.

Theorem 7. *Let $j > 1$. The longest prefix of the Fibonacci infinite word that is an abelian repetition of period F_j has length $F_j(F_{j+1}+F_{j-1}+1)-2$ if j is even or $F_j(F_{j+1}+F_{j-1})-2$ if j is odd.*

Proof. (Sketch) It is enough to prove that the length of the head and of the tail can be maximal (i.e., the head and the tail both have length $F_j - 1$) in the case of the maximal exponent described in Proposition 6. Indeed, by Proposition 4 and Theorem 6, there exists exactly one factor v that has a Parikh vector different from the Parikh vector $\mathcal{Q}$ of all the other factors, including the factor that is the prefix of f of length F_j. The corresponding interval of this unique factor is either $L_0(\alpha, F_j)$ or $L_{F_j}(\alpha, F_j)$, and therefore it is not the interval $\phi - 1$ belongs to.

Therefore, the Parikh vector of the prefix of length F_j of f is $\mathcal{Q}$ and the Parikh vector of the prefix of length $F_j - 1$ of f is $\mathcal{P} \subset \mathcal{Q}$. This proves that the head has maximal length. Since the exponent considered in Proposition 6 is maximal, the factor of length F_j must be the unique factor v with a different Parikh vector. But, by geometric considerations, the prefix of v of length $|v| - 1 = F_j - 1$ has other different factors of the same length that share its Parikh vector $\mathcal{P}$. These other factors are prefixes of at least one factor of length F_j that has Parikh vector $\mathcal{Q}$. Therefore, $\mathcal{P} \subset \mathcal{Q}$. This proves that also the tail has maximal length. □

Corollary 4. *Let $j > 1$ and let k_j be the maximal exponent of a prefix of the Fibonacci word that is an abelian repetition of period F_j. Then*

$$\lim_{j \to \infty} \frac{k_j}{F_j} = \sqrt{5}.$$

Proof. We proved in a previous theorem that over a bigger set of repetitions the superior limit tends to $\sqrt{5}$. Therefore, in this case, it must be smaller than or equal to $\sqrt{5}$. The equality follows from the fact that $\frac{F_{j+1}+F_{j-1}}{F_j}$ converges to $\sqrt{5}$. Since the sequence $\frac{F_{j+1}}{F_j}$ converges to ϕ and the sequence $\frac{F_{j-1}}{F_j}$ converges to $\phi - 1$, then the sequence

$$\frac{F_{j+1} + F_{j-1}}{F_j} = \frac{F_{j+1}}{F_j} + \frac{F_{j-1}}{F_j}$$

converges to

$$\phi + \phi - 1 = \frac{\sqrt{5}+1}{2} + \frac{\sqrt{5}-1}{2} = \sqrt{5}.$$

□



Fig. 5. Longest abelian repetition of period m that is a prefix of the Fibonacci word for $m = 2, 3, 5$. (a) For $m = 2$, the longest abelian repetition has length $8 = 1 + 3p + 1$. (b) For $m = 3$, the longest abelian repetition has length $19 = 2 + 5p + 2$. (c) For $m = 5$, the longest abelian repetition has length $58 = 4 + 10p + 4$.

j	2	3	4	5	6	7	8	9	10	11
F_j	2	3	5	8	13	21	34	55	89	144
$lp(F_j)$	8	19	58	142	388	985	2616	6763	17798	46366
$\lvert\sqrt{5} - k_j/F_j\rvert * 10^2$	23.6	12.5	8.393	1.732	5.98	0.25	2.69	0.037	1.087	0.005

Table 1. The length of the longest prefix ($lp(F_j)$) of the Fibonacci word having abelian period F_j for $j = 2, \ldots, 11$. The table also reports rounded distances (multiplied by 10^2) between $\sqrt{5}$ and the ratio between the exponent $k_j lp(F_j)/F_j$ of the longest prefix of the Fibonacci word having abelian period F_j and F_j (see Corollary 4).

In Fig. 5, we give a graphical representation of the longest prefix of the Fibonacci infinite word that is an abelian repetition of period m for $m = 2, 3$ and 5.

In Table 1, we give the length $lp(F_j)$ of the longest prefix of the Fibonacci infinite word that is an abelian repetition of period F_j, for $j = 2, \ldots, 11$, computed using the formula of Theorem 7. We also show the values of the distance between $\sqrt{5}$ and the ratio between the maximal exponent of a prefix of the Fibonacci infinite word having abelian period F_j and F_j.

Recall that the Fibonacci (finite) words are defined by $f_0 = \mathsf{b}$, $f_1 = \mathsf{a}$, and for every $j > 1$, $f_{j+1} = f_j f_{j-1}$. So, for every j, one has $|f_j| = F_j$.

As a consequence of the formula given in Theorem 7, we have the following result on the smallest abelian periods of the Fibonacci words.

Theorem 8. *For $j \geq 3$, the (smallest) abelian period of the word f_j is the n-th Fibonacci number F_n, where $n = \lfloor j/2 \rfloor$ if $j = 0, 1, 2 \mod 4$, or $n = 1 + \lfloor j/2 \rfloor$ if $j = 3 \mod 4$.*

Proof. (Sketch) It is possible to prove that no abelian repetition that is a prefix of the Fibonacci infinite word of period m, with $F_n < m < F_{n+1}$, $n > 2$, has length greater than the length of the abelian repetition that is a prefix of the Fibonacci infinite word of period F_n. Indeed, if m is such that $F_n < m < F_{n+1}$, $n > 2$, one has that $\{m(\phi - 1)\}$

is not too close to an integer and then it is possible to use the same technique developed in the proof of Theorem 1. By using Proposition 7, it is sufficient to find the smallest index n such that $|f_j| = F_j$ is greater than the length of the longest abelian repetition in the Fibonacci infinite word of period F_{n-1}. A computation that makes use of Theorem 2 gives the desired formula. □

For example, the abelian period $f_4 =$ **abaab** is $2 = F_2 = \lfloor 4/2 \rfloor$, since one can write $f_4 =$ **a** · **ba** · **ab**; the abelian period of $f_5 =$ **abaababa** is $2 = F_2$; the abelian period of $f_6 =$ **abaababaabaab** is $3 = F_3$; the abelian period of $f_7 =$ **abaababaabaababaababa** is $5 = F_4$.

In Table 2 we report the abelian periods of the first Fibonacci words.

j	3	4	5	6	7	8	9	10	11	12	13	14	15	16
a. p. of f_j	F_2	F_2	F_2	F_3	F_4	F_4	F_4	F_5	F_6	F_6	F_6	F_7	F_8	F_8

Table 2. The (smallest) abelian periods of the Fibonacci words f_j for $j = 3, \ldots, 16$.

References

1. A. Aho. Algorithms for Finding Patterns in Strings. In J. van Leeuwen, editor, *Handbook of Theoret. Comput. Sci.*, pages 257–300. Elsevier Science Publishers B. V., Amsterdam, the Netherlands, 1990.
2. S. Avgustinovich, A. Glen, B. Halldórsson, and S. Kitaev. On shortest crucial words avoiding abelian powers. *Discrete Appl. Math.*, 158(6):605–607, 2010.
3. S. Avgustinovich, J. Karhumäki, and S. Puzynina. On abelian versions of Critical Factorization Theorem. *RAIRO Theor. Inform. Appl.*, 46:3–15, 2012.
4. J. Berstel. Sturmian and episturmian words (a survey of some recent results). In S. Bozapalidis and G. Rohonis, editors, *CAI 2007*, volume 4728 of *Lecture Notes in Comput. Sci.*, pages 23–47. Springer, 2007.
5. J. Berstel, A. Lauve, C. Reutenauer, and F. Saliola. *Combinatorics on Words: Christoffel Words and Repetition in Words*, volume 27 of *CRM monograph series*. American Mathematical Society, 2008.
6. F. Blanchet-Sadri, J. I. Kim, R. Mercas, W. Severa, and S. Simmons. Abelian square-free partial words. In *Proceedings of the 4th International Conference Language and Automata Theory and Applications*, volume 6031 of *Lecture Notes in Comput. Sci.*, pages 94–105. Springer, 2010.
7. F. Blanchet-Sadri, A. Tebbe, and A. Veprauskas. Fine and Wilf's theorem for abelian periods in partial words. In *Proceedings of the 13th Mons Theoret. Comput. Sci. Days*, 2010.
8. M. Bucci, A. De Luca, and L. Zamboni. Some characterizations of Sturmian words in terms of the lexicographic order. *Fundamenta Informaticae*, 116(1-4):25–33, 2012.
9. P. Burcsi, F. Cicalese, G. Fici, and Zs. Lipták. Algorithms for jumbled pattern matching in strings. *Int. J. Found. Comput. Sci.*, 23(2):357–374, 2012.
10. J. Cassaigne, G. Richomme, K. Saari, and L. Zamboni. Avoiding Abelian powers in binary words with bounded Abelian complexity. *Int. J. Found. Comput. Sci.*, 22(4):905–920, 2011.
11. M. Christou, M. Crochemore, and C. S. Iliopoulos. Identifying all abelian periods of a string in quadratic time and relevant problems. *CoRR*, abs/1207.1307, 2012.

12. F. Cicalese, G. Fici, and Zs. Lipták. Searching for Jumbled Patterns in Strings. In *Proceedings of the Prague Stringology Conference 2009, PSC 2009*, pages 105–117. Czech Technical University in Prague, 2009.
13. S. Constantinescu and L. Ilie. Fine and Wilf's theorem for abelian periods. *Bull. Eur. Assoc. Theoret. Comput. Sci. EATCS*, 89:167–170, 2006.
14. M. Crochemore, L. Ilie, and W. Rytter. Repetitions in strings: Algorithms and combinatorics. *Theoret. Comput. Sci.*, 410(50):5227–5235, 2009.
15. L. J. Cummings and W. F. Smyth. Weak repetitions in strings. *J. Combin. Math. Combin. Comput.*, 24:33–48, 1997.
16. J. Currie and N. Rampersad. Fixed points avoiding Abelian k-powers. *Journal of Combinatorial Theory, Series A*, 119(5):942–948, 2012.
17. J. D. Currie and A. Aberkane. A cyclic binary morphism avoiding Abelian fourth powers. *Theoret. Comput. Sci.*, 410(1):44–52, 2009.
18. F. Dekking. Strongly non-repetitive sequences and progression-free sets. *Journal of Combinatorial Theory, Series A*, 27(2):181–185, 1979.
19. M. Domaratzki and N. Rampersad. Abelian primitive words. In *Proceedings of the 15th Conference on Developments in Language Theory*, volume 6795 of *Lecture Notes in Comput. Sci.*, pages 204–215. Springer, 2011.
20. P. Erdös. Some unsolved problems. *Magyar Tud. Akad. Mat. Kutato. Int. Kozl.*, 6:221–254, 1961.
21. A. Glen, J. Justin, and G. Pirillo. Characterizations of finite and infinite episturmian words via lexicographic orderings. *European Journal of Combinatorics*, 29(1):45–58, 2008.
22. G. H. Hardy and E. M. Wright. *An Introduction to the Theory of Numbers.* Clarendon Press, Oxford, 1979. 5th edition.
23. M. Huova, J. Karhumäki, A. Saarela, and K. Saari. Local squares, periodicity and finite automata. In *Rainbow of Computer Science*, pages 90–101, 2011.
24. C. S. Iliopoulos, D. Moore, and W. F. Smyth. A Characterization of the Squares in a Fibonacci String. *Theoret. Comput. Sci.*, 172(1-2):281–291, 1997.
25. O. Jenkinson and L. Q. Zamboni. Characterisations of balanced words via orderings. *Theoret. Comput. Sci.*, 310(1-3):247–271, 2004.
26. V. Keränen. Abelian squares are avoidable on 4 letters. In *Proceedings of the 19th International Colloquium on Automata, Languages and Programming*, volume 623 of *Lecture Notes in Comput. Sci.*, pages 41–52. Springer-Verlag, 1992.
27. R. Klette and A. Rosenfeld. Digital straightness: a review. *Discrete Appl. Math.*, 139(1-3):197–230, Apr. 2004.
28. R. Kolpakov and G. Kucherov. Finding Maximal Repetitions in a Word in Linear Time. In *Proceedings of the 40th Annual Symposium on Foundations of Computer Science*, FOCS '99, pages 596–604. IEEE Computer Society, 1999.
29. M. Lothaire. *Algebraic Combinatorics on Words.* Cambridge University Press, Cambridge, U.K., 2002.
30. A. Mateescu, A. Salomaa, K. Salomaa, and S. Yu. A sharpening of the parikh mapping. *RAIRO Theor. Inform. Appl.*, 35(6):551–564, 2001.
31. F. Mignosi. Infinite Words with Linear Subword Complexity. *Theoret. Comput. Sci.*, 65(2):221–242, 1989.
32. F. Mignosi. On the number of factors of Sturmian words. *Theoret. Comput. Sci.*, 82:71–84, 1991.
33. F. Mignosi and G. Pirillo. Repetitions in the Fibonacci infinite word. *RAIRO Theor. Inform. Appl.*, 26:199–204, 1992.
34. F. Mignosi and A. Restivo. Characteristic Sturmian words are extremal for the critical factorization theorem. *Theoret. Comput. Sci.*, 454(0):199 – 205, 2012.
35. R. J. Parikh. On context-free languages. *J. Assoc. Comput. Mach.*, 13(4):570–581, 1966.
36. D. Perrin and A. Restivo. A note on Sturmian words. *Theoret. Comput. Sci.*, 429:265–272, 2012.
37. S. Puzynina and L. Q. Zamboni. Abelian returns in Sturmian words. In *Proceedings 8th International Conference Words, WORDS 2011*, volume 63 of *EPTCS*, pages 179–188, 2011.

38. N. Pytheas Fogg. *Substitutions in Dynamics, Arithmetics and Combinatorics*, volume 1794 of *Lecture Notes in Math.* Springer, 2002.
39. G. Richomme, K. Saari, and L. Zamboni. Balance and Abelian complexity of the Tribonacci word. *Advances in Applied Mathematics*, 45(2):212–231, 2010.
40. G. Richomme, K. Saari, and L. Zamboni. Abelian complexity of minimal subshifts. *Journal of the London Mathematical Society*, 83(1):79–95, 2011.
41. A. Salomaa. Subword histories and associated matrices. *Theoret. Comput. Sci.*, 407(1-3):250–257, 2008.
42. A. Samsonov and A. Shur. On Abelian repetition threshold. *RAIRO Theor. Inform. Appl.*, 46:147–163, 2012.